

# PHYSICAL ACTIVITY AND PSYCHOLOGICAL WELL-BEING IN THE ADULT POPULATION DURING THE COVID-19 PANDEMIC: A SYSTEMATIC REVIEW

*by* Hanna Oktasya Ross

---

**Submission date:** 25-Jul-2023 11:41AM (UTC+0700)

**Submission ID:** 2136455169

**File name:** Revisi\_Hanna\_2\_1\_1.docx (1.02M)

**Word count:** 7515

**Character count:** 47185



## PHYSICAL ACTIVITY AND PSYCHOLOGICAL WELL-BEING IN THE ADULT POPULATION DURING THE COVID-19 PANDEMIC: A SYSTEMATIC REVIEW

anna Oktasya Ross<sup>1\*</sup>, Erico Pratama Nugraha<sup>2</sup>, Ardhito Faza Akhnaf<sup>3</sup>, Nanum Sofia<sup>4</sup>

<sup>1,2,3,4</sup> Department of Psychology, Faculty of Psychology and Socio-Cultural Sciences, Universitas Islam Indonesia, Yogyakarta

\*Correspondence Author: [hannaoktasya@gmail.com](mailto:hannaoktasya@gmail.com)

### ARTICLE INFO

#### Article History:

Received : January 19, 2023

Accepted : July 14, 2023

Published : July 31, 2023

#### DOI:

<https://doi.org/10.26553/jikm.2023.14.2.127-135>

#### Available online at

<http://ejournal.fkm.unsri.ac.id/index.php/jikm>

### ABSTRACT

The COVID-19 pandemic has become a global crisis that has caused various impacts on the psychological well-being and physical activity of the community. Correspondingly, physical activity is known to have a relationship and affect psychological well-being. Therefore, this study aimed to obtain a comprehensive picture of the level and impact of the activity on psychological well-being in the adult population as well as recommendations for physical activity that could be done to improve psychological well-being during the COVID-19 pandemic. The research method used was a systematic literature review with the PRISMA (Preferred Reporting Item for Systematic Review and Meta-Analytic) analysis method. Search literature on the EBSCOhost, ProQuest, and ScienceDirect databases. Literature searches were limited to research articles published between January 1, 2020, to May 20, 2021. The initial search yielded 5,136 articles and finally 15 articles that met the criteria. The results of the review revealed that during the COVID-19 pandemic, there was a decrease in the level of physical activity and psychological well-being, especially in the adult population, especially women and young adult groups. In addition, it was found that physical activity was either directly or indirectly associated with symptoms of depression, stress, anxiety, and positive feelings and emotions. Physical activity with moderate intensity has the most significant influence on psychological well-being. The forms of activity included lifting weights, yoga, double tennis, aerobics, muscle fitness, and explosive fitness. Based on this, it is certain that physical activity with moderate intensity has the most significant impact on psychological well-being, especially in increasing positive feelings and emotions and reducing depression, stress, and anxiety in adults.

**Keywords:** adult, COVID-19 pandemic, physical activity, psychological well-being

## Introduction

Since the end of 2019, the COVID-19 pandemic has evolved into a global crisis. COVID-19 is a respiratory infection illness characterized by SARS-CoV-2/severe acute respiratory syndrome coronavirus-2.<sup>1</sup> The most common symptoms of COVID-19 infection are fever, coughing, sneezing, and shortness of breath.<sup>2</sup> The risk of contracting COVID-19 increases if a person has comorbid hypertension, diabetes mellitus, is an active smoker,<sup>3,4</sup> has close contact with a COVID-19 patient, and has a history of travel to COVID-19-infected areas.<sup>5</sup> The virus was first suspected of spreading in the Huanan fish market in Wuhan, China,<sup>2</sup> and less than a month after being reported; the disease spread rapidly in many provinces in China, South Korea, Japan, and Thailand.<sup>6</sup> Then, from February to April 2021, globally confirmed cases of COVID-19 increased significantly, increasing by more than 50% from 2,486,744 confirmed cases to 5,740,164 confirmed cases of death, increasing by more than 11% from 67,645 to 93,854 deaths.<sup>7</sup>

Several new policies have been implemented to prevent the spread of COVID-19 and reduce the positive number. Quarantine, isolation,<sup>8-11</sup> lockdown,<sup>12-14</sup> and work from home are among the new policies.<sup>15,16</sup> This policy alters people's social habits by requiring them to always wash their hands, wear masks when leaving the house, maintain a safe distance when interacting with others, avoid crowds, reduce activities outside the home and mobility from one location to another, and even work and go to school from home online.<sup>17</sup> This policy also has an influence on people's and society's psychological well-being, including the development of Post-Traumatic Stress Disorder (PTSD), frustration, confusion, anxiety, fear of affection, sleep disturbances, and feelings of helplessness. Even if the condition worsens, the excessive fear of infection from COVID-19 can lead to xenophobia and suicidal thoughts.<sup>18</sup>

In line with this, it has been reported that there was an increase in psychological disorders during the COVID-19 pandemic, particularly among adults.<sup>19,20</sup> Adults are also confirmed to be the most vulnerable group in several countries to COVID-19 infection.<sup>21,22</sup> These conditions can also disrupt a person's psychological well-being, potentially causing a variety of problems ranging from depression to anxiety,<sup>23,24</sup> reduced intensity of relationships with other people,<sup>25</sup> loss of purpose in life,<sup>26</sup> burnout severity,<sup>27,28</sup> inhibiting personal growth,<sup>29</sup> social isolation, loneliness,<sup>30</sup> and self-harm.<sup>31</sup> Some of these consequences occur in adults,<sup>18</sup> particularly in cases of depression and anxiety.<sup>32</sup>

In general, good psychological well-being in adults will increase body immunity, productivity, and motivation to do activities and work, resulting in greater stability in dealing with various situations.<sup>33</sup> Meanwhile, good psychological well-being in adults, particularly during the COVID-19 pandemic, will result in the ability to respond to stress by engaging in good self-coping behaviors.<sup>34</sup> It is supported by numerous studies on psychological well-being during the COVID-19 pandemic. Based on search results from five international database portals, there are 14,673 studies

discussing psychological well-being during the COVID-19 pandemic. This indirectly demonstrates that psychological well-being is a global issue that requires further investigation.

Psychological well-being is defined as the ability of a person's psychological aspects to function normally and positively.<sup>35</sup> A person's psychological well-being is influenced by a variety of factors, encompassing gender,<sup>36,37</sup> age,<sup>38</sup> social support, optimism,<sup>39,40</sup> gratitude,<sup>41</sup> emotional control,<sup>42</sup> goal attainment,<sup>43</sup> and physical activity.<sup>43-46</sup> Physical activity is one of the psychological well-being factors that will be discussed in this study. The researcher selected physical activity as a factor because it was one of the topics studied extensively during the COVID-19 pandemic, particularly in relation to psychological well-being. A total of 2,072 studies discussing physical activity and psychological well-being during the COVID-19 pandemic were discovered using search results from five international database portals. Several studies have discovered a connection between physical activity and psychological well-being.<sup>47-49</sup> Furthermore, physical activity helps to reduce levels of anxiety, and depression,<sup>50-52</sup> sleep disturbances,<sup>53</sup> enhances cognitive performance,<sup>54</sup> and has a positive effect on the physique.<sup>50-52,55</sup>

Based on the preceding explanation, the objective of this study is to obtain a comprehensive picture of the level and impact of physical activity on psychological well-being in adults. This is consistent with previous research, which indicates that the adult population is the most vulnerable to COVID-19 infection and is psychologically affected. This study was conducted through a systematic review of several previous studies because many studies explained the level and impact of physical activity on psychological well-being during the COVID-19 pandemic. However, no comprehensive research has been conducted in the adult population. As a result, it is expected that this study will provide an overview to the general public, particularly the adult population, of the level and impact of physical activity on psychological well-being, as well as recommendations for what physical activity can be implemented to enhance psychological well-being during the COVID-19 pandemic.

The research problem formulation encompasses: 1) What is the level and impact of physical activity on psychological well-being in the adult population during the COVID-19 pandemic? 2) What forms of physical activity can be done during the COVID-19 pandemic to improve psychological well-being?

## Methods

This research procedure started with determining research topics, formulating research questions, identifying keywords, searching and identifying databases, and finally, reading and analyzing. The method employed in this study was a literature study with a systematic literature review approach, which was a systematic literature review method that identifies, evaluates, and interprets all findings on a research topic to answer predetermined research questions. Literature

searches were limited to research articles published between January 1, 2020, to May 20, 2021. Article searches were conducted online employing the words “physical activity” OR “physical exercise” AND “psychological well-being” OR “mental health” OR “mental illness ” AND “COVID-19” in titles, keywords, and abstracts in journal articles contained in the ProQuest, ScienceDirect, and EBSCOhost portal databases. The literature search process started by reviewing all search results' titles, keywords, and abstracts, filtering and classifying them based on predetermined criteria, then the final process of the review process was carried out by experts. In addition, the data extraction process was also carried out by reviewing a number of selected articles and writing down the main findings. Then the extraction results would be synthesized by grouping them into findings that could answer the objectives of this study.

The analytical method administered was the Preferred Reporting Item for Systematic Review and Meta-Analytic (PRISMA) method. All articles that passed the selection were then reviewed and summarized based on predetermined criteria starting from the objective, author's name, year of publication, number of respondents, instruments used, and research results. The criteria for this study were demonstrated in Table 1.

**Table 1. Inclusion and Exclusion Criteria**

Category	Inclusion Criteria	Exclusion Criteria
Publication Criteria	Empirical research articles (research articles) and peer-reviewed in full-text form Research from any country and written in English or Indonesian language.	Non-empirical research, literature review studies, meta-analyses, case studies, reports, proceedings, book chapters, and essays.
Period of time	Articles published from January 1, 2020 to May 20, 2021.	
Study Content	Articles containing physical activity or physical exercise, psychological well-being or mental health, and the COVID-19 pandemic	Subjective well-being, spiritual well-being, physical training, physical health, physical abilities, long covid, measures/ scale
Population	Adult age group (18 – 60 years)	Babies, children, teenagers, and the elderly

The initial search produced 5,132 articles from three databases: science direct (977 articles), ProQuest (4,094 articles), and EBSCOhost (65 articles). The PRISMA method was employed in the flow of the article selection process (Figure 1). Then, the first search concentrates only on articles that contain the keywords "physical activity" OR "physical exercise" AND "psychological well-being" OR "mental health" OR "mental illness" AND "COVID-19". Afterward, duplicate articles were removed until only 4,595 remained. The following stage, screening, was successful in excluding 3,384 articles, as only 1,211 fulfilled the criteria. The screening criteria were as follows: 1) Articles published between January 1, 2020 and May 20, 2021, 2) Full-text, peer-reviewed articles in academic journals or research papers, 3) Articles in Indonesian and English; 4) Open access to articles. Then, at the eligibility stage, the 1,121 articles were screened by reading their titles, abstracts, and keywords. In the end, 241 articles were obtained, while 970 other articles were excluded because they did not fulfill the established criteria, particularly regarding the fact that not all of the articles contained the main topic, as well as the keywords that had been planned at the

beginning. Furthermore, 214 articles were excluded at the included stage since the research subjects were not of adult age. Thus, 27 articles out of a total of 241 articles fulfilled the criteria. Then, from the 27 articles, a final stage of screening was performed, which included reading the articles in their entirety, and 15 articles were identified as meeting the criteria and being designated as material for a systematic review of this research.




Figure 1. Article Selection Process Flowchart

## Results

The majority of the 15 articles mentioned above utilized quantitative research with a cross-sectional study design. Various studies have been conducted in Europe, America, Asia, and Australia. Participants for the study were obtained through online surveys on social media, websites, email, mass media, and government media. The instrument employed in the research was a questionnaire that assessed physical activity (IPAQ, MET, GLTEQ, BREQ-3, AAS), psychological well-being (PGWBI, GHQ-12, WHO Five Well-Being Index (WHO-5), MHC-SF), depression (CES-D, PHQ-9, BDI-II, DASS-21), anxiety (GAD-7, STAI, HADS, DASS-21), stress (PSS, DASS-21, IES-6), sleep quality (PSQI, Sleep Quality Survey), mood (POMS), positive and negative emotions (HOT). Table 2 summarizes the main characteristics of the eligible studies. Table 3 illustrates the definitions, measurements, and results.

Table 2. Study of Main Characteristics

Author	Country	Study Design	Population	Sample	Gender	Age Range	Media
Althumiri et al, 2021 <sup>57</sup>	Saudi Arabia	Cross-Sectional Study	Public Society	n = 8333	P = 4.192 L = 4.141	> 18 years	Online survey
Cecchini et al., 2021 <sup>58</sup>	Spain	Longitudinal Study	Public Society	n = 595	P = 342 L = 253	18-84 years	Online survey, social media & mass media
Denget al., 2020 <sup>59</sup>	China	17 Cross-Sectional Study	University Students	n = 1607	P = 566 L = 1.041	18-22 years	Online survey
Dun et al, 2021 <sup>60</sup>	China	Cross-Sectional and Retrospective Study	University Students	n = 12.880	P = 10.340 L = 2.540	18-21 years	Online survey
Green et al, 2021 <sup>47</sup>	17 countries	Cross-Sectional Study	University Students	n = 7.303	P = 6.129 L = 1.147 Blank = 27	Median 47 years	Online survey
Kilani et al., 2020 <sup>61</sup>	Iraq and	Cross-Sectional Study	University Students	n = 1723	P = 806 L = 917	18 – 65 years	E-mail and social media
Lesser et al., 2020 <sup>62</sup>	Canada	Cross-Sectional Study	University Students	n = 1098	P = 871 L = 215	> 19 years	Social media, mass, govern ment
Lin, et al, 2020 <sup>63</sup>	China	17 Cross-Sectional Study	University Students	n = 628	P = 405 L = 220	Average 20 years	Online survey
Marashi, et al 2020 <sup>64</sup>	Canada & other countries	Cross-Sectional Study	Public Society	n = 1669	P = 1218 L = 249 Blank = 11	> 18 years	Online survey
Maugeri et al., 2020 <sup>65</sup>	Italy	Cross-Sectional Study	Public Society	n = .524	P = 1.426 L = 1.098	21-60 years	Online survey, social media, and e-mail
Puccinelli et al., 2021 <sup>66</sup>	Brazil	Cross-Sectional Study	Public Society	n = 1853	P = 1.110 L = 743	> 18 years	Online surveys, websites, e-mails, and social media
Qin et al., 2020 <sup>67</sup>	China	Cross-Sectional Study	Public Society	n = 12.107	P = 6.474 L = 5.366	18-80 years	Online survey
Reigal et al., 2021 <sup>68</sup>	Spain	Correlational Studies	Public Society	n = 328	P = 209 L = 119	19-59 years	Online survey
Stanton et al., 2020 <sup>69</sup>	Australia	Cross-Sectional Study	Public Society	n = 1491	P = 999 L = 484	> 18 years	Online survey, social media, and e-mail
Zhang et al., 2020 <sup>70</sup>	China	Longitudinal Study	University Students	n = 66	P = 42 L = 25	20-21 years	Online survey

Table 3.Objectives, Instruments, and Outcomes

Author	Objective	Measurement/ Instrument	Result
Alhumiri et al., 2021 <sup>57</sup>	Exploring the relationship between physical activity and risk and symptoms of depression and anxiety in the Saudi population	PHQ-9, GAD-7	Moderate physical activity (such as light weight lifting, yoga or relaxation, and doubles tennis) can improve general mental health, especially by reducing levels of anxiety and depression. Meanwhile, high physical activity (weight lifting, jogging, aerobics, and fast cycling) can reduce symptoms of depression, but not anxiety.
Cecchini et al., 2021 <sup>58</sup>	Assessing the prevalence of depressive symptoms, physical activity and providing guidelines for effective and efficient physical activity during isolation during the Covid pandemic in Spain	IPAQ, MET, Six Item Self-reported questionnaire by Kandel and Davies (1982)	Depressive symptoms increased significantly during isolation during the pandemic, particularly in women and the younger-age group. Physical activity has a negative relationship with depressive symptoms. Starting with moderate physical activity (4 hours/week) can reduce symptoms of mild depression and continue with high physical activity (16 hours/week) can further reduce symptoms of depression significantly.
Denget al., 2020 <sup>59</sup>	Determining the relationship between mental health and lifestyle associated with the physical activity of students in Wuhan and evaluating a website-based physical education program	DASS-21, Sport-Related Lifestyle Scale	The majority of participants had normal mental health status, and only a few participants experienced depression, anxiety, and stress. Significant correlation between depression, anxiety, and stress with regular exercise 1-2 times a week for an adequate duration of > 1 hour. Meanwhile, web-based sports can be employed as an alternative, but among students, it is considered less effective/unsatisfactory.
Dun et al., 2021 <sup>60</sup>	Assessing the prevalence of depression in young adults during the COVID-19 pandemic and exploring its relationship to different types of physical fitness	BDI-II & IPAQ-LC	Moderate and high physical activity demonstrate a lower risk of depression than participants with low levels of physical activity in which men experience a more significant decrease in physical activity. Moderate and high levels of aerobic, explosive, and muscular fitness can reduce the risk of depression.
Green et al., 2021 <sup>47</sup>	Examining the differences and relationships between mental health and concerns associated with COVID-19 and exploring the effects of meditation and physical activity on mental health	CDC, SRHI, PSS, IES-6, HADS	Anxiety levels, stress symptoms, depression, and PTSD have increased more severely, particularly in people in countries with high cases of COVID-19 and strict social distancing restrictions. Physical activity and meditation are significant mediators in reducing symptoms of stress, depression, PTSD, and anxiety, particularly those related to concerns about COVID-19
Kilani et al., 2020 <sup>61</sup>	Identifying the extent to which lifestyle behaviors such as physical activity, sleep, and diet contribute to mental well-being during the COVID-19 pandemic	WHO Five Well-Being Index (WHO-5), FFQ, PSQI, IPAQ	Physical activity is the primary predictor of better mental health status. The other predictors of improved mental well-being are quality eating and sleeping patterns. Mental well-being scores were higher among participants with moderate and high levels of physical activity (PA). The male population has significantly higher levels of physical activity and mental well-being.
Lesser & Nienhuis., 2020 <sup>62</sup>	Investigating the impact of the pandemic on the physical activity and psychological well-being of	GLTEQ, BREQ-3, MHC-SF, NRS, GAD-7	There was a decrease in physical activity in which participants who were more active maintained a physical level and connectedness to nature and demonstrated lower levels of activity anxiety, and higher social, emotional, psychological well-being, autonomy, and



Author	Objective	Measurement/ Instrument	Result
Lin et al., 2020 <sup>63</sup>	Canadians and exploring the differences between outdoor physical activity and exposure to nature based on classifications of anxiety and general well-being Examining the relationship between depressive symptoms and physical activity among Chinese college students and examining the effect of gender roles on that relationship during the COVID-19 pandemic. Investigating changes in physical activity, sedentary behavior, and mental health before and during the COVID-19 pandemic	IPAQ-SF, CES-D	self-motivation than participants who were inactive Participants who conduct the moderate physical activity with other people possess higher mental health than those who perform physical activity alone. However, it did not occur in participants who did heavy and light physical activities. A third of participants illustrate increased rates of depression and decreased levels of physical activity during the COVID-19 pandemic. Gender with high masculinity traits have lower rates of depression. Physical activity has a positive correlation with depression levels, particularly moderate physical activity can reduce depression levels.
Marashi et al., 2020 <sup>64</sup>	Investigating changes in physical activity, sedentary behavior, and mental health before and during the COVID-19 pandemic	PASB-Q GAD-7 PHQ-9 PSS	During the pandemic, sedentary activity (sitting) decreased, psychological stress increased, and also symptoms of anxiety and depression increased, particularly in the group with low physical activity. Participants were not motivated to perform physical activity because they were too anxious, lack of social support, or limited access to equipment and facilities. The young adult age group (18-29 years) illustrates higher levels of depression and anxiety than other groups, also experiencing a decrease in physical activity and an increase in sedentary activity.
Maugeri et al., 2020 <sup>65</sup>	Examining changes in physical activity levels and physical well-being of individuals during self-quarantine in Italy	IPAQ & PGWBI	Physical activity during the pandemic has decreased significantly in both adult women and men. High anxiety, depressed mood, positive well-being, self-control, especially in the group of women and young adults.
Puccinelli et al., 2021 <sup>66</sup>	Identifying the impact of social distancing on changes in physical activity levels and their relationship to mood (depression and anxiety) as well as people's total income during the Covid pandemic in Spain, Brazil	IPAQ, PHQ-9, GAD-7	During social distancing, physical activity decreased significantly, and 30% of participants reported moderate and high levels of depression and anxiety. Mood (depressive and anxiety symptoms) is negatively correlated with physical activity and family income and is associated with the younger age group.
Qin et al., 2020 <sup>67</sup>	Identifying the impact of the COVID-19 pandemic on the level and lifestyle characteristics of the Chinese population during quarantine	PANAS, IPAQ-SF	It was reported that there was a decrease in physical activity and an increase in the ration of screen time, especially for males. Individuals with vigorous physical activity have better emotional states and less screen time than those with light physical activity.
Reigal et al., 2021 <sup>68</sup>	Identifying the relationship between physical activity, mood conditions, and self-assessment of health during	IPAQ, POMS, STAI, GHQ-12	There is a positive relationship between physical activity and emotional state. Higher positive emotions were discovered in males and were associated with the effects of high physical activity. Physical activity is positively associated with perceptions of health, anxiety, and mood (anger, depression, enthusiasm, tension). Moderate physical activity predicts better mood and health perceptions than vigorous

Author	Objective	Measurement/ Instrument	Result
Stanton et al., 2020 <sup>69</sup>	Investigating the relationship between psychological disorders and behavioral changes in physical activity, sleep, smoking, and alcohol use during social isolation in Australia.	DASS-21, AUDIT-C, AAS, Sleep Quality Survey	physical activity and positively affects levels of anger, depression, enthusiasm, stress, and anxiety. Men and younger age groups (19 -39 years) demonstrate better psychological health when performing high/intense physical activity. On the other hand, women and older groups (40 -59 years) illustrated better psychological health when conducting moderate physical activity.
Zhang et al., 2020 <sup>70</sup>	Identifying the impact and relationship between the COVID-19 pandemic and mental health (depression, anxiety, stress, sleep quality), and exploring the effects of appropriate mitigation/physical activity.	IPAQ-SF, DASS 21, PSQI, BPAQ,	The COVID-19 pandemic has significantly reduced the quality of sleep for young adults and has implications for increasing negative emotions, especially stress, and anxiety. Women experienced a more significant decrease in physical activity. The death rate and severity of the COVID-19 outbreak have an indirect effect on negative emotions (stress, depression, and anxiety), sleep quality and reduce people's level of aggressiveness, in which the majority of participants, particularly women, felt more anxious about COVID-19, stress, anxiety, and depression in themselves. Moderate physical activity significantly reduces negative emotions.

According to the findings of a review of 15 articles, the most common symptoms of psychological disorders during the COVID-19 pandemic were depression, anxiety, stress, sleep disturbances, and various emotional and mood disorders demonstrated in Table 4.

**Table 4. Psychological Well-Being**

Category	Author
Depression	Althumiri et al, 2021 <sup>57</sup> ; Cecchini et al., 2021 <sup>58</sup> ; Deng et al., 2020 <sup>59,60</sup> ; Dun et al, 2021 <sup>60</sup> ; Green et al, 2021 <sup>47</sup> ; Liu et al, 2020 <sup>63</sup> ; Marashi et al, 2020 <sup>64</sup> ; Stanton et al., 2020 <sup>69</sup> ; Puccinelli et al., 2021 <sup>66</sup> ; Reigal et al, 2021 <sup>68</sup> ; Zhang et al., 2020 <sup>70</sup>
Anxiety	Althumiri et al, 2021 <sup>57</sup> ; Cecchini et al., 2021 <sup>58</sup> ; Deng et al., 2020 <sup>59</sup> ; Dun et al, 2021 <sup>60</sup> ; Green et al, 2021 <sup>47</sup> ; Lin et al, 2020 <sup>63</sup> ; Marashi et al, 2020 <sup>64</sup> ; Stanton et al., 2020 <sup>69</sup> ; Puccinelli et al., 2021 <sup>66</sup> ; Reigal et al, 2021 <sup>68</sup> ; Zhang et al., 2020 <sup>70</sup>
Stress	Deng et al., 2020 <sup>59</sup> ; Green et al, 2021 <sup>47</sup> ; Marashi et al, 2020 <sup>64</sup> ; Stanton et al., 2020 <sup>69</sup> ; Zhang et al., 2020 <sup>70</sup>
Emotional and Mood Disorders	Lesser & Nienhuis., 2020 <sup>62</sup> ; Maugeri et al., 2020 <sup>65</sup> ; Qin et al., 2020 <sup>67</sup> ; Reigal et al, 2021 <sup>68</sup>
Sleep Disorders	Kilani et al., 2020 <sup>61</sup> ; Stanton et al., 2020 <sup>69</sup> ; Zhang et al., 2020 <sup>70</sup>

The primary question addressed in this study is the level and impact of physical activity on psychological well-being in the adult population during the COVID-19 pandemic. In accordance with the review's findings, the majority of studies discovered that the amount of physical activity in the adult population had decreased.<sup>63-69</sup> On the other hand, sedentary physical activities, such as sitting, lying down, and using screens, have elevated during the COVID-19 pandemic.<sup>64,67</sup> This is also indirectly due to the policy of limiting activities outside the home during the COVID-19 pandemic, such as quarantine,<sup>65,67</sup> isolation,<sup>58,69</sup> lockdown,<sup>68</sup> work from home,<sup>15,16</sup> and social distancing.<sup>66</sup> Furthermore, the subject is less motivated to engage in physical activities because they are too anxious, lack social support, or are ineffective and have limited access to physical activity equipment and facilities.<sup>59,64</sup>

Both men and women experience a decrease in physical activity,<sup>63,65,66</sup> with men experiencing a greater decrease than women.<sup>67</sup> Other studies, however, have revealed that women are more likely than men to experience a decrease in physical activity.<sup>60,61,70</sup> Meanwhile, the adult population, particularly those aged 18-24 years, experienced a more significant decrease in physical activity than other individuals.<sup>64,67</sup>

Based on the results of a systematic review that has been carried out, it shows that the level of physical activity either directly or indirectly also impacts psychological well-being. In general, doing physical activity can improve mental health,<sup>61,62,65</sup> self-control,<sup>65</sup> reduce symptoms of depression, anxiety, and stress,<sup>47,64,66,68,69</sup> PTSD,<sup>47</sup> and negative mood.<sup>65,68</sup> Meanwhile, based explicitly on intensity, the review results show that moderate-intensity physical activity has the

most significant effect on psychological well-being compared to low and high-intensity physical activity. These results are also supported by several studies which report that moderate-intensity physical activity can reduce symptoms of depression,<sup>57,58,60,63</sup> anxiety,<sup>57</sup> negative emotions, and feelings.<sup>70,68</sup> However, the results of other studies also report that physical activity with high or vigorous intensity is effective only on some psychological well-being, such as reducing symptoms of depression,<sup>57,58,60</sup> anxiety,<sup>65</sup> and increasing positive emotions and moods.<sup>67,68</sup>

Then, respond to the second problem formulation, which is about physical activity recommendations that can be performed <sup>19</sup> during the COVID-19 pandemic to enhance <sup>6</sup> psychological well-being. In accordance with the findings of a systematic review, the most effective form of physical activity for enhancing psychological well-being is to engage in moderate-intensity physical activity 1-2 times per week for at least 1 hour <sup>59</sup> or approximately 4 hours per week.<sup>58</sup> Performing light weights, yoga, doubles tennis,<sup>57</sup> aerobics, muscular fitness (push-ups, sit-ups, squats, dance, digging, shovelling, climbing stairs), and explosive fitness (high steps, long jumps, and repeated squat jumps) are all recommended forms of moderate-intensity physical activity.<sup>60</sup> Furthermore, forms of high-intensity physical activity such as weight lifting, jogging, aerobics, cycling, muscular, and explosive fitness for a long period of time are recommended because they possess an effect on some psychological well-being. <sup>60</sup>

### Discussion

Some research reveals that women are more likely than males to experience a decline in physical activity.<sup>60,61,70</sup> However, physical activity decreases for both men and women,<sup>63,65,66</sup> with men reporting a higher reduction than women.<sup>67</sup> Meanwhile, the adult population, particularly those aged 18-24 years, suffered a greater reduction in physical activity than the general population.<sup>64,67</sup> These findings are justified by other investigations that age and gender have a large role in dropping the number of physical activities, and even this effect tends to raise the risk of mental health.<sup>71</sup> According to the findings of a comprehensive review, the degree of physical activity has an impact on psychological well-being, either directly or indirectly. Physical activity, in general, can promote mental health<sup>61,62,65</sup> and reduce symptoms of anxiety and stress.<sup>47,64,66,68,69</sup> Furthermore, these findings are confirmed by the study that shows the level of physical activity has a crucial effect on enhancing mental health and reducing the probability of depression.<sup>72</sup>

<sup>2</sup> The study's major question is the level and influence of physical exercise on psychological well-being in the adult population during the COVID-19 pandemic. According to the review's findings, the majority of studies discovered that physical activity levels in the adult population had declined.<sup>63-69</sup> On the other hand, Sedentary physical behaviors, including sitting, lying down, and utilizing screens, have increased. <sup>1</sup> During the COVID-19 pandemic.<sup>64,67</sup> These findings are confirmed by research conducted in Italy, during the COVID-19 pandemic, the vast majority of

Italian can not do anything. Because of that, sedentary activities have increased, whereas physical activity has decreased.<sup>73</sup> Furthermore, the victim of the pandemic is less motivated to participate in physical activities because they are very nervous, lack social support, are ineffective, and have restricted access to physical exercise equipment and facilities.<sup>59,64</sup> These findings are related to another research that has the same findings. It reveals that the poor state of emotions during the COVID-19 pandemic leads us to obtain less motivation in order to do physical activities.<sup>74</sup>

Meanwhile, the review results reveal that moderate-intensity physical activity has the greatest influence on psychological well-being when compared to low and high-intensity physical activity. Furthermore, moderate-intensity physical activity has been shown to lower symptoms of depression,<sup>57,58,60,63</sup> anxiety,<sup>57</sup> negative emotions, and sentiments.<sup>70,68</sup> The findings of other studies indicate that high-intensity physical activity is useful only on particular aspects of psychological well-being, such as reducing symptoms of depression,<sup>57,58,60</sup> anxiety,<sup>65</sup> and raising positive feelings and moods.<sup>67,68</sup> These findings are supported by some research that reveals the effect of intense physical activity on psychological well-being. The study justifies that the intensity of physical activity, such as week-intensity and vigorous-intensity, has a substantial role in achieving psychological well-being.<sup>73</sup>

Then, respond to the second problem formulation, which is about physical activity recommendations that can be performed during the COVID-19 pandemic to enhance psychological well-being. According to the findings of a comprehensive study, the most effective form of physical activity for improving psychological well-being is moderate-intensity physical activity 1-2 times per week for at least 1 hour or about 4 hours each week.<sup>58</sup> That suggestion has the same main point as other physical activity recommendations during the COVID-19 pandemic.

Moderate-intensity physical exercise is defined as lifting light weights, yoga, doubles tennis,<sup>57</sup> aerobics, muscular fitness (push-ups, sit-ups, squats, dance, digging, shoveling, climbing stairs), and explosive fitness (high steps, long jumps, and repeated squat jumps).<sup>60</sup> Furthermore, long-term high-intensity physical activity such as weight lifting, jogging, aerobics, cycling, muscular and explosive fitness are encouraged since they have an effect on some psychological well-being.<sup>60</sup> That kind of recommendation has been sported by other studies that show some physical exercise which has the advantage of enhancing the immune system and reducing the risk of diseases during the COVID-19 pandemic.<sup>76</sup>

### Conclusion

Based on the findings and discussion, it is possible to conclude that the adult population's level of physical activity and psychological well-being decreased during the COVID-19 pandemic, particularly among women and young adults. Physical activity could have an impact on psychological well-being both directly and indirectly, particularly in increasing positive feelings

and emotions <sup>3</sup> and decreasing symptoms of depression, stress, and anxiety in the adult population. Furthermore, regular moderate-intensity physical activity 1-2 times per week for more than 1 hour, or approximately 4 hours per week, produced the most significant effect on psychological well-being. Performing light weights, yoga, doubles tennis, aerobics, muscular fitness, and explosive fitness are several moderate-intensity physical activities that are recommended. Furthermore, forms of high-intensity physical activity such as weight lifting, jogging, aerobics, cycling, muscular fitness, and explosive fitness of strong duration are also highly recommended as they are also effective in several psychological well-being.

### **Acknowledgment**

Not applicable.

### **Funding**

During and after this research was conducted, there was no funding for the research.

### **Conflict of Interest**

During and after this research was conducted, there was no conflict of interest.

### **References**

1. WHO. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. <https://www.who.int/director-general/speeches/detail/who-director-general-opening-remarks-at-the-media-briefing-on-COVID-19--11-march-2020>. 2020.
2. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun.* 2020 May;109:102433. <https://doi.org/10.1016/j.jaut.2020.102433>
3. Cai H. Sex difference and smoking predisposition in patients with COVID-19. *Lancet Respir Med.* 2020 Apr;8(4):e20. [https://doi.org/10.1016/S2213-2600\(20\)30117-X42](https://doi.org/10.1016/S2213-2600(20)30117-X42)
4. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med.* 2020 Apr;8(4):e21. [https://doi.org/10.1016/S2213-2600\(20\)30116-8](https://doi.org/10.1016/S2213-2600(20)30116-8)
5. CDC. Interim US guidance for risk assessment and public health management of persons with potential coronavirus disease 2019 (COVID-19) exposures: geographic risk and contacts of laboratory-confirmed cases. <https://stacks.cdc.gov/view/cdc/85723>. 2020.

6. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*. 2020 Feb;395(10223):497–506. [https://doi.org/10.1016/s0140-6736\(20\)30183-5](https://doi.org/10.1016/s0140-6736(20)30183-5)
7. WHO. WHO Coronavirus (COVID-19) Dashboard. <https://covid19.who.int/>. 2021.
8. Aronna MS, Guglielmi R, Moschen LM. A model for COVID-19 with isolation, quarantine and testing as control measures. *Epidemics*. 2021 Mar;34:100437. <https://doi.org/10.1016/j.epidem.2021.100437>
9. Baldacchino G. Extra-territorial quarantine in pandemic times. *Polit Geogr*. 2021 Mar;85:102302. <https://doi.org/10.1016/j.polgeo.2020.102302>
10. Gondim JAM, Machado L. Optimal quarantine strategies for the COVID-19 pandemic in a population with a discrete age structure. *Chaos Solitons Fractals*. 2020 Nov;140:110166. <https://doi.org/10.1016/j.chaos.2020.110166>
11. Wang X, Wang J, Shen J, Ji JS, Pan L, Liu H, et al. Facilities for Centralized Isolation and Quarantine for the Observation and Treatment of Patients with COVID-19. *Engineering*. 2021 Jul;7(7):908–13. <https://doi.org/10.1016/j.eng.2021.03.010>
12. Coccia M. The relation between length of lockdown, numbers of infected people and deaths of COVID-19, and economic growth of countries: Lessons learned to cope with future pandemics similar to COVID-19 and to constrain the deterioration of economic system. *Science of The Total Environment*. 2021 Jun;775:145801. <https://doi.org/10.1016/j.scitotenv.2021.145801>
13. Fu Y, Jin H, Xiang H, Wang N. Optimal lockdown policy for vaccination during COVID-19 pandemic. *Financ Res Lett*. 2022 Mar;45:102123. <https://doi.org/10.1016/j.frl.2021.102123>
14. Jia Z, Xu S, Zhang Z, Cheng Z, Han H, Xu H, et al. Association between mental health and community support in lockdown communities during the COVID-19 pandemic: Evidence from rural China. *J Rural Stud*. 2021 Feb;82:87–97. <https://doi.org/10.1016/j.jrurstud.2021.01.015>
15. Alipour JV, Fadinger H, Schymik J. My home is my castle – The benefits of working from home during a pandemic crisis. *J Public Econ*. 2021 Apr;196:104373. <https://doi.org/10.1016/j.jpubeco.2021.104373>
16. M V, Mehroliya S, Alagarsamy S, Balachandran AK. Work from home in the pandemic era: Loss of mental equilibrium? *Asian J Psychiatr*. 2021 Jan;55:102490. <https://doi.org/10.1016/j.ajp.2020.102490>
17. Yanuarita HA, Haryati S. Pengaruh COVID-19 terhadap kondisi sosial budaya di kota malang dan konsep strategis dalam penanganannya. *Jurnal Ilmiah Widya Sosiopolitika*. 2021 Jan 20;2(2):58. <https://doi.org/10.24843/JIWSWP.2020.v02.i02.p01>

18. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet*. 2020 Mar;395(10227):912–20. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
19. Knolle F, Ronan L, Murray GK. The impact of the COVID-19 pandemic on mental health in the general population: a comparison between Germany and the UK. *BMC Psychol*. 2021 Dec 24;9(1):60. <https://doi.org/10.1186/s40359-021-00565-y>
20. PSDKJI. Masalah psikologis terkait pandemi COVID-19 di Indonesia. <https://pdskji.org/home>. 2020.
21. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*. 2020 Mar;395(10229):1054–62. [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3)
22. Asirvatham ES, Sarman CJ, Saravanamurthy SP, Mahalingam P, Maduraipandian S, Lakshmanan J. Who is dying from COVID-19 and when? An Analysis of fatalities in Tamil Nadu, India. *Clin Epidemiol Glob Health*. 2021 Jan;9:275–9. <https://doi.org/10.1016/j.cegh.2020.09.010>
23. Liu Y, Yue S, Hu X, Zhu J, Wu Z, Wang J, et al. Associations between feelings/behaviors during COVID-19 pandemic lockdown and depression/anxiety after lockdown in a sample of Chinese children and adolescents. *J Affect Disord*. 2021 Apr;284:98–103. <https://doi.org/10.1016/j.jad.2021.02.001>
24. Luo M, Guo L, Yu M, Jiang W, Wang H. The psychological and mental impact of coronavirus disease 2019 (COVID-19) on medical staff and general public – A systematic review and meta-analysis. *Psychiatry Res*. 2020 Sep;291:113190. <https://doi.org/10.1016/j.psychres.2020.113190>
25. Sachser C, Olaru G, Pfeiffer E, Brähler E, Clemens V, Rassenhofer M, et al. The immediate impact of lockdown measures on mental health and couples' relationships during the COVID-19 pandemic - results of a representative population survey in Germany. *Soc Sci Med*. 2021 Jun;278:113954. <https://doi.org/10.1016/j.socscimed.2021.113954>
26. Na PJ, Tsai J, Hill ML, Nichter B, Norman SB, Southwick SM, et al. Prevalence, risk and protective factors associated with suicidal ideation during the COVID-19 pandemic in U.S. military veterans with pre-existing psychiatric conditions. *J Psychiatr Res*. 2021 May;137:351–9. <https://doi.org/10.1016/j.jpsychires.2021.03.021>
27. Cooper K, Hards E, Moltrecht B, Reynolds S, Shum A, McElroy E, et al. Loneliness, social relationships, and mental health in adolescents during the COVID-19 pandemic. *J Affect Disord*. 2021 Jun;289:98–104. <https://doi.org/10.1016/j.jad.2021.04.016>



28. Mion G, Hamann P, Saleten M, Plaud B, Baillard C. Psychological impact of the COVID-19 pandemic and burnout severity in French residents: A national study. *Eur J Psychiatry*. 2021 Jul;35(3):173–80. <https://doi.org/10.1016/j.ejpsy.2021.03.005>
29. Asmundson GJG, Paluszek MM, Taylor S. Real versus illusory personal growth in response to COVID-19 pandemic stressors. *J Anxiety Disord*. 2021 Jun;81:102418. <https://doi.org/10.1016/j.janxdis.2021.102418>
30. Williams CYK, Townson AT, Kapur M, Ferreira AF, Nunn R, Galante J, et al. Interventions to reduce social isolation and loneliness during COVID-19 physical distancing measures: A rapid systematic review. *PLoS One*. 2021 Feb 17;16(2):e0247139. <https://doi.org/10.1371/journal.pone.0247139>
31. Hawton K, Lascelles K, Brand F, Casey D, Bale L, Ness J, et al. Self-harm and the COVID-19 pandemic: A study of factors contributing to self-harm during lockdown restrictions. *J Psychiatr Res*. 2021 May;137:437–43. <https://doi.org/10.1016/j.jpsychires.2021.03.028>
32. Panchal N, Kamal R, Cox C, Garfield R. The Implications of COVID-19 for Mental Health and Substance Use. <https://www.kff.org/coronavirus-COVID-19/issue-brief/the-implications-of-COVID-19-for-mental-health-and-substance-use/>. 2021.
33. Khairunisa NS, Safitri DR, Angelia D, Taufan M, Sihalo ED. Produktivitas dan depresi di Indonesia: Analisis data Indonesia Family Life Survey 2014. *Jurnal Ekonomi Pembangunan*. 2019 Dec 31;27(2):75–84. <https://doi.org/10.14203/JEP.27.2.2019.75-84>
34. Tuason MaT, Güss CD, Boyd L. Thriving during COVID-19: Predictors of psychological well-being and ways of coping. *PLoS One*. 2021 Mar 15;16(3):e0248591. <https://doi.org/10.1371/journal.pone.0248591>
35. Sumakul Y, Ruata S. Kesejahteraan psikologis dalam masa pandemi COVID-19. *Journal of Psychology" Humanlight"*. 2020;1(1):1–7.
36. Brebabama A, Listyandini RA. Gambaran tingkat kesejahteraan psikologis penyandang tunanetra dewasa muda. *Mediapsi*. 2016 Jun 1;02(01):1–10. <https://doi.org/10.21776/ub.mps.2016.002.01.1>
37. Izzati UA, Mulyana OP. Perbedaan jenis kelamin dan status pernikahan dalam kesejahteraan psikologis guru. *Psychocentrum Review*. 2021 Mar 25;3(1):63–71. <https://doi.org/10.26539/pcr.31567>
38. Lee SH, Choi I, Choi E, Lee M, Kwon Y, Oh B, et al. Psychological well-being and gene expression in Korean adults: The role of age. *Psychoneuroendocrinology*. 2020 Oct;120:104785. <https://doi.org/10.1016/j.psyneuen.2020.104785>

39. Khoirunnisa A, Ratnaningsih IZ. Optimisme dan kesejahteraan psikologis pada mahasiswa pendidikan dokter Fakultas Kedokteran Universitas Diponegoro. *Jurnal EMPATI*. 2016 Jan 27;5(1):1–4. <https://doi.org/10.14710/empati.2016.14936>
40. Safarina NA, Munir A, Nuraini N. Hubungan Harga Diri dan Optimisme dengan Kesejahteraan Subjektif pada Mahasiswa Magister Psikologi Universitas Medan Area. *Tabularasa: Jurnal Ilmiah Magister Psikologi*. 2019;1(1):39–48. <https://doi.org/10.31289/analitika.v8i2.868>
41. Aisyah A, Chisol R. Rasa syukur kaitannya dengan kesejahteraan psikologis pada Guru Honorer Sekolah Dasar. *Proyeksi*. 2020 Sep 26;13(2):109. <http://dx.doi.org/10.30659/jp.13.2.109-122>
42. Konradus N, Harsanti I. Peranan Kecerdasan Emosi dan Kesejahteraan Psikologis Terhadap Kepuasan Kerja Guru pada Sebuah Yayasan Pendidikan Islam di Bekasi. *Jurnal Psikologi*. 2017;8(2).
43. Verner-Filion J, Vallerand RJ, Amiot CE, Mocuano I. The two roads from passion to sport performance and psychological well-being: The mediating role of need satisfaction, deliberate practice, and achievement goals. *Psychol Sport Exerc*. 2017 May;30:19–29. <https://doi.org/10.1016/j.psychsport.2017.01.009>
44. Kim J, Lee S, Chun S, Han A, Heo J. The effects of leisure-time physical activity for optimism, life satisfaction, psychological well-being, and positive affect among older adults with loneliness. *Annals of Leisure Research*. 2017 Aug 8;20(4):406–15. <https://doi.org/10.1080/11745398.2016.1238308>
45. Abidin S, Welch RK, Byron-Daniel J, Meyrick J. The effectiveness of physical activity interventions in improving well-being across office-based workplace settings: a systematic review. *Public Health*. 2018 Jul;160:70–6. <https://doi.org/10.1016/j.puhe.2018.03.029>
46. Humphreys L, Crank H, Frith G, Speake H, Reece LJ. Bright spots, physical activity investments that work: Active Everyday, Sheffield’s physical activity service for all people living with and beyond cancer. *Br J Sports Med*. 2019 Jul;53(13):837–8. <https://doi.org/10.1136/bjsports-2017-098672>
47. Green J, Huberty J, Puzia M, Stecher C. The Effect of Meditation and Physical Activity on the Mental Health Impact of COVID-19–Related Stress and Attention to News Among Mobile App Users in the United States: Cross-sectional Survey. *JMIR Ment Health*. 2021 Apr 13;8(4):e28479. <https://doi.org/10.2196/28479>
48. Maugeri G, Musumeci G. Adapted Physical Activity to Ensure the Physical and Psychological Well-Being of COVID-19 Patients. *J Funct Morphol Kinesiol*. 2021 Jan 29;6(1):13. <https://doi.org/10.1016/j.heliyon.2020.e04315>

49. Nienhuis CP, Lesser IA. The Impact of COVID-19 on Women's Physical Activity Behavior and Mental Well-Being. *Int J Environ Res Public Health*. 2020 Dec 4;17(23):9036. <https://doi.org/10.3390/ijerph17239036>
50. Blough J, Loprinzi PD. Experimentally investigating the joint effects of physical activity and sedentary behavior on depression and anxiety: A randomized controlled trial. *J Affect Disord*. 2018 Oct;239:258–68. <https://doi.org/10.1016/j.jad.2018.07.019>
51. Carter T, Pascoe M, Bastounis A, Morres ID, Callaghan P, Parker AG. The effect of physical activity on anxiety in children and young people: a systematic review and meta-analysis. *J Affect Disord*. 2021 Apr;285:10–21. <https://doi.org/10.1016/j.jad.2021.02.026>
52. Forte C, McDowell C, MacDonncha C, Herring MP. Depressive symptoms differ across Physical Activity Status based on comorbid anxiety and depression status among adolescents. *Ment Health Phys Act*. 2020 Oct;19:100338. <https://doi.org/10.1016/j.mhpa.2020.100338>
53. Duncan MJ, Holliday EG, Oftedal S, Buman M, Brown WJ. Joint association of physical activity and sleep difficulties with the incidence of hypertension in mid-age Australian women. *Maturitas*. 2021 Jul;149:1–7. <https://doi.org/10.1016/j.maturitas.2021.04.006>
54. Sousa CV, Hwang J, Cabrera-Perez R, Fernandez A, Misawa A, Newhook K, et al. Active video games in fully immersive virtual reality elicit moderate-to-vigorous physical activity and improve cognitive performance in sedentary college students. *J Sport Health Sci*. 2022 Mar;11(2):164–71. <https://doi.org/10.1016/j.jshs.2021.05.002>
55. Woods JA, Hutchinson NT, Powers SK, Roberts WO, Gomez-Cabrera MC, Radak Z, et al. The COVID-19 pandemic and physical activity. *Sports Medicine and Health Science*. 2020 Jun;2(2):55–64. <https://doi.org/10.1016/j.smhs.2020.05.006>
56. Mengist W, Soromessa T, Legese G. Method for conducting systematic literature review and meta-analysis for environmental science research. *MethodsX*. 2020 Jan 1;7:100777. <https://doi.org/10.1016/j.mex.2019.100777>
57. Althumiri NA, Basyouni MH, BinDhim NF. Exploring the Association Between Physical Activity and Risk of Mental Health Disorders in Saudi Arabian Adults: Cross-sectional Study. *JMIR Public Health Surveill*. 2021 Apr 14;7(4):e25438. <https://doi.org/10.2196/25438>
58. Cecchini JA, Carriedo A, Fernández-Río J, Méndez-Giménez A, González C, Sánchez-Martínez B, et al. A longitudinal study on depressive symptoms and physical activity during the Spanish lockdown. *International Journal of Clinical and Health Psychology*. 2021 Jan;21(1):100200. <https://doi.org/10.1016/j.ijchp.2020.09.001>
59. Deng CH, Wang JQ, Zhu LM, Liu HW, Guo Y, Peng XH, et al. Association of Web-Based Physical Education With Mental Health of College Students in Wuhan During the COVID-

- 19 Outbreak: Cross-Sectional Survey Study. *J Med Internet Res.* 2020 Oct 5;22(10):e21301. <https://doi.org/10.2196/21301>
60. Dun Y, Ripley-Gonzalez JW, Zhou N, Li Q, Chen M, Hu Z, et al. The association between prior physical fitness and depression in young adults during the COVID-19 pandemic—a cross-sectional, retrospective study. *PeerJ.* 2021 Mar 11;9:e11091.
61. Kilani HA, Bataineh MF, Al-Nawayseh A, Atiyat K, Obeid O, Abu-Hilal MM, et al. Healthy lifestyle behaviors are major predictors of mental wellbeing during COVID-19 pandemic confinement: A study on adult Arabs in higher educational institutions. *PLoS One.* 2020 Dec 14;15(12):e0243524. <https://doi.org/10.7717/peerj.11091>
62. Lesser IA, Nienhuis CP. The Impact of COVID-19 on Physical Activity Behavior and Well-Being of Canadians. *Int J Environ Res Public Health.* 2020 May 31;17(11):3899. <https://doi.org/10.3390/ijerph17113899>
63. Lin J, Guo T, Becker B, Yu Q, Chen ST, Brendon S, et al. Depression is Associated with Moderate-Intensity Physical Activity Among College Students During the COVID-19 Pandemic: Differs by Activity Level, Gender and Gender Role. *Psychol Res Behav Manag.* 2020 Dec; Volume 13:1123–34. <https://doi.org/10.2147/PRBM.S277435>
64. Marashi MY, Nicholson E, Ogrodnik M, Fenesi B, Heisz JJ. A mental health paradox: Mental health was both a motivator and barrier to physical activity during the COVID-19 pandemic. *PLoS One.* 2021 Apr 1;16(4):e0239244. <https://doi.org/10.1371/journal.pone.0239244>
65. Maugeri G, Castrogiovanni P, Battaglia G, Pippi R, D'Agata V, Palma A, et al. The impact of physical activity on psychological health during COVID-19 pandemic in Italy. *Heliyon.* 2020 Jun;6(6):e04315. <https://doi.org/10.1016/j.heliyon.2020.e04315>
66. Puccinelli PJ, da Costa TS, Seffrin A, de Lira CAB, Vancini RL, Nikolaidis PT, et al. Reduced level of physical activity during COVID-19 pandemic is associated with depression and anxiety levels: an internet-based survey. *BMC Public Health.* 2021 Dec 1;21(1):425. <https://doi.org/http://dx.doi.org/10.1186/s12889-021-10470-z>
67. Qin F, Song Y, Nassis GP, Zhao L, Dong Y, Zhao C, et al. Physical activity, screen time, and emotional well-being during the 2019 Novel Coronavirus outbreak in China. *Int J Environ Res Public Health.* 2020 Jul 17;17(14):5170. <https://doi.org/http://dx.doi.org/10.3390/ijerph17145170>
68. Reigal RE, Páez-Maldonado JA, Pastrana-Brincones JL, Morillo-Baro JP, Hernández-Mendo A, Morales-Sánchez V. Physical Activity Is Related to Mood States, Anxiety State and Self-Rated Health in COVID-19 Lockdown. *Sustainability.* 2021 May 13;13(10):5444. <https://doi.org/10.3390/su13105444>

69. Stanton R, To QG, Khalesi S, Williams SL, Alley SJ, Thwaite TL, et al. Depression, Anxiety and Stress during COVID-19: Associations with Changes in Physical Activity, Sleep, Tobacco and Alcohol Use in Australian Adults. *Int J Environ Res Public Health*. 2020 Jun 7;17(11):4065. <https://doi.org/http://dx.doi.org/10.3390/ijerph17114065>
70. Zhang Y, Zhang H, Ma X, Di Q. Mental Health Problems during the COVID-19 Pandemics and the Mitigation Effects of Exercise: A Longitudinal Study of College Students in China. *Int J Environ Res Public Health*. 2020 May 25;17(10):3722. <https://doi.org/http://dx.doi.org/10.3390/ijerph17103722>
71. Pieh C, Budimir S, Probst T. The effect of age, gender, income, work, and physical activity on mental health during coronavirus disease (COVID-19) lockdown in Austria. *Journal of psychosomatic research*. 2020 Sep 1;136:110186. <https://doi.org/10.1016/j.jpsychores.2020.110186>
72. Rebar AL, Stanton R, Geard D, Short C, Duncan MJ, Vandelanotte C. A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health psychology review*. 2015 Aug 7;9(3):366-78. <https://doi.org/10.1080/17437199.2015.1022901>
73. Maugeri G, Castrogiovanni P, Battaglia G, Pippi R, D'Agata V, Palma A, Di Rosa M, Musumeci G. The impact of physical activity on psychological health during COVID-19 pandemic in Italy. *Heliyon*. 2020 Jun 1;6(6). <https://doi.org/10.1016/j.heliyon.2020.e04315>
74. Füzéki E, Groneberg DA, Banzer W. Physical activity during COVID-19 induced lockdown: recommendations. *Journal of Occupational Medicine and Toxicology*. 2020 Dec;15(1):1-5. <https://doi.org/10.1186/s12995-020-00278-9>
75. Polero P, Rebollo-Seco C, Adsuar JC, Pérez-Gómez J, Rojo-Ramos J, Manzano-Redondo F, Garcia-Gordillo MÁ, Carlos-Vivas J. Physical activity recommendations during COVID-19: narrative review. *International journal of environmental research and public health*. 2021 Jan;18(1):65. <https://doi.org/10.3390/ijerph18010065>
76. Chagas EF, Biteli P, Candeloro BM, Rodrigues MA, Rodrigues PH. Physical exercise and COVID-19: a summary of the recommendations. *Aims Bioengineering*. 2020;7(4):236-41.

# PHYSICAL ACTIVITY AND PSYCHOLOGICAL WELL-BEING IN THE ADULT POPULATION DURING THE COVID-19 PANDEMIC: A SYSTEMATIC REVIEW

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

14%

INTERNET SOURCES

15%

PUBLICATIONS

5%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://covid19dataportal.es">covid19dataportal.es</a> Internet Source	1%
2	<a href="http://www.wjgnet.com">www.wjgnet.com</a> Internet Source	1%
3	<a href="https://journals.plos.org">journals.plos.org</a> Internet Source	1%
4	<a href="https://link.springer.com">link.springer.com</a> Internet Source	1%
5	Min Luo, Lixia Guo, Mingzhou Yu, Haiyan Wang. "The psychological and mental impact of coronavirus disease 2019 (COVID-19) on medical staff and general public – A systematic review and meta-analysis", <i>Psychiatry Research</i> , 2020 Publication	1%
6	Grazia Maugeri, Paola Castrogiovanni, Giuseppe Battaglia, Roberto Pippi et al. "The impact of physical activity on psychological	1%

# health during Covid-19 pandemic in Italy", Heliyon, 2020

Publication

---

7	wjgnet.com Internet Source	1 %
8	www.mdpi.com Internet Source	1 %
9	www.researchgate.net Internet Source	<1 %
10	Marc Lochbaum, Thaís Zanatta, Zişan Kazak. "The 2 × 2 Achievement Goals in Sport and Physical Activity Contexts: A Meta-Analytic Test of Context, Gender, Culture, and Socioeconomic Status Differences and Analysis of Motivations, Regulations, Affect, Effort, and Physical Activity Correlates", European Journal of Investigation in Health, Psychology and Education, 2019 Publication	<1 %
11	pubcovid19.pt Internet Source	<1 %
12	Submitted to Universitas Mercu Buana Student Paper	<1 %
13	Submitted to University of Plymouth Student Paper	<1 %
14	ijssr.ridwaninstitute.co.id	

---

Internet Source

<1 %

15

[m.amedeo.com](https://m.amedeo.com)

Internet Source

<1 %

16

[www.minervamedica.it](http://www.minervamedica.it)

Internet Source

<1 %

17

[www.scienceexcel.com](http://www.scienceexcel.com)

Internet Source

<1 %

18

[mdpi-res.com](https://mdpi-res.com)

Internet Source

<1 %

19

[www.cambridge.org](http://www.cambridge.org)

Internet Source

<1 %

20

Sunil Luthra, Shruti Agrawal, Anil Kumar, Manu Sharma, Sudhanshu Joshi, Jatin Kumar. "Psychological well-being of young adults during COVID-19 pandemic: Lesson learned and future research agenda", Heliyon, 2023

Publication

<1 %

21

Submitted to Universitas Islam Indonesia

Student Paper

<1 %

22

M.S. Aronna, R. Guglielmi, L.M. Moschen. "A model for COVID-19 with isolation, quarantine and testing as control measures", Epidemics, 2021

Publication

<1 %



23	<a href="http://hdl.handle.net">hdl.handle.net</a> Internet Source	<1 %
24	<a href="http://media.neliti.com">media.neliti.com</a> Internet Source	<1 %
25	<a href="http://www.pegegog.net">www.pegegog.net</a> Internet Source	<1 %
26	<a href="http://9pdf.net">9pdf.net</a> Internet Source	<1 %
27	Rafael E. Reigal, José A. Páez-Maldonado, José L. Pastrana-Brincones, Juan P. Morillo-Baro et al. "Physical Activity Is Related to Mood States, Anxiety State and Self-Rated Health in COVID-19 Lockdown", Sustainability, 2021 Publication	<1 %
28	Submitted to University of Kentucky Student Paper	<1 %
29	<a href="http://ejournal.unesa.ac.id">ejournal.unesa.ac.id</a> Internet Source	<1 %
30	<a href="http://www.thejmch.com">www.thejmch.com</a> Internet Source	<1 %
31	Cana Aksoy Poyraz, Büşra Uçar Bostan, Elif Burcu Ersungur Çelik, Beril Kara Çelik. "The impact of COVID-19 pandemic and lockdown on emergency psychiatry department	<1 %

# admissions", Research Square Platform LLC, 2023

Publication

---

32	<a href="https://diposit.ub.edu">diposit.ub.edu</a> Internet Source	<1 %
33	<a href="https://peerj.com">peerj.com</a> Internet Source	<1 %
34	<a href="https://publichealth.jmir.org">publichealth.jmir.org</a> Internet Source	<1 %
35	<a href="https://staging-publichealth.jmir.org">staging-publichealth.jmir.org</a> Internet Source	<1 %
36	<a href="https://doaj.org">doaj.org</a> Internet Source	<1 %
37	<a href="https://journals.lww.com">journals.lww.com</a> Internet Source	<1 %
38	<a href="https://oss.jomh.org">oss.jomh.org</a> Internet Source	<1 %
39	<a href="https://www.aimspress.com">www.aimspress.com</a> Internet Source	<1 %
40	<a href="https://www.elsevier.es">www.elsevier.es</a> Internet Source	<1 %
41	<a href="https://www.researchsquare.com">www.researchsquare.com</a> Internet Source	<1 %
42	<a href="https://www.revistapsiquiatria.pt">www.revistapsiquiatria.pt</a> Internet Source	<1 %

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On